

REFLECTIONS, BENDINGS, AND PENTAGONS

SASHA ANAN'IN

ABSTRACT. We study relations between reflections in (positive or negative) points in the complex hyperbolic plane. It is easy to see that the reflections in the points q_1, q_2 obtained from p_1, p_2 by moving p_1, p_2 along the geodesic generated by p_1, p_2 and keeping the (dis)distance between p_1, p_2 satisfy the *bending relation* $R(q_2)R(q_1) = R(p_2)R(p_1)$. We show that a generic isometry $F \in \mathrm{SU}(2, 1)$ is a product of 3 reflections, $F = R(p_3)R(p_2)R(p_1)$, and describe all such decompositions: two decompositions are connected by finitely many bendings involving $p_1, p_2/p_2, p_3$ and geometrically equal decompositions differ by an isometry centralizing F .

Any relation between reflections gives rise to a representation $H_n \rightarrow \mathrm{PU}(2, 1)$ of the *hyperelliptic group* H_n generated by $r_1, \dots, r_n$ with the defining relations $r_n \dots r_1 = 1, r_j^2 = 1$. The theorem mentioned above is essential to the study of the Teichmüller space $\mathcal{TH}_n$. We describe all nontrivial representations of H_5 , called *pentagons*, and conjecture that they are faithful and discrete.

What do you feed the people generated by reflections?

— A. A. Kirillov to E. B. Vinberg on the way to Baikal Lake

1. Introduction

We will deal with the complex hyperbolic plane $\mathbb{H}_{\mathbb{C}}^2$. In our notation, we follow [AGr] and [AGG]. Let V denote a 3-dimensional $\mathbb{C}$ -linear space equipped with a hermitian form $\langle -, - \rangle$ of signature $++-$. By definition, $\mathbb{H}_{\mathbb{C}}^2 := \mathrm{B}V$ and $\partial\mathbb{H}_{\mathbb{C}}^2 := \mathrm{S}V$, where

$$\mathrm{B}V := \{p \in \mathbb{P}_{\mathbb{C}}V \mid \langle p, p \rangle < 0\}, \quad \mathrm{S}V := \{p \in \mathbb{P}_{\mathbb{C}}V \mid \langle p, p \rangle = 0\}, \quad \mathrm{E}V := \{p \in \mathbb{P}_{\mathbb{C}}V \mid \langle p, p \rangle > 0\}$$

(see [AGr] and [AGG] for details).

Let us start with some heuristic considerations. For many geometries, it is well known that every isometry is a product of involutions. It is curious to understand to what extent such a decomposition is unique. In other words, we study the ‘defining relations’ $R_n \dots R_2 R_1 = 1$ and $R_j^2 = 1$.

Consider $\mathrm{PU}V$, the group of holomorphic isometries of $\mathbb{H}_{\mathbb{C}}^2$. Every involution in $\mathrm{PU}V$ is a reflection $R(p)$ in some nonisotropic point $p \in \mathbb{P}_{\mathbb{C}}V \setminus \mathrm{S}V$ (positive or negative).

- The relation $R(p)R(p) = 1$ is said to be a *cancellation*.

Clearly, these are the only relations with $n = 2$.

- For pairwise orthogonal $p_1, p_2, p_3 \in \mathbb{P}_{\mathbb{C}}V \setminus \mathrm{S}V$, the reflections $R(p_1), R(p_2), R(p_3)$ commute and $R(p_3)R(p_2)R(p_1) = 1$. We call these relations *orthogonal*.

It is easy to see that these are the only relations with $n = 3$ (see Remark 3.2).

Suppose that distinct points $p_1, p_2 \in \mathbb{P}_{\mathbb{C}}V \setminus \mathrm{S}V$ generate a hyperbolic¹ projective line $L(p_1, p_2)$. Then the polar point p to $L(p_1, p_2)$ is positive and $R(p_2)R(p_1) = R(p_2)R(p)R(p)R(p_1) = R(p'_2)R(p'_1)$, where

¹2000 *Mathematics Subject Classification*. 30F60 (51M10, 57S30).

¹ $L(p_1, p_2)$ is *hyperbolic, spherical, euclidean* if the signature of $\mathbb{C}p_1 + \mathbb{C}p_2$ is respectively $+-, ++, +0$.

$p'_1, p'_2 \in L(p_1, p_2)$ are respectively orthogonal to p_1, p_2 . So, with the use of cancellation and orthogonal relations, we can simultaneously alter the signs of p_1, p_2 if the projective line $L(p_1, p_2)$ is hyperbolic.

Given distinct and nonorthogonal points $p_1, p_2 \in \mathbb{P}_{\mathbb{C}}V \setminus SV$, we can move p_1, p_2 along the geodesic $G \setminus \{p_1, p_2\}$ and keep constant the distance (tance) between them. The new points q_1, q_2 satisfy the relation $R(q_2)R(q_1) = R(p_2)R(p_1)$. More formally: There exists a one-parameter subgroup $B : (\mathbb{R}, +) \rightarrow \text{PU}V$ such that $B(s)$ centralizes $R(p_2)R(p_1)$, $\text{dist}(p_1, B(s)p_1) = \text{dist}(p_1, p_2)|s|$, $B(1)p_1 = p_2$ if p_1, p_2 have the same sign, and $B(1)p_1 \in L(p_1, p_2)$ is orthogonal to p_2 if p_1, p_2 have different signs. For $q_1 := B(s)p_1$ and $q_2 := B(s)p_2$, we have $R(q_2)R(q_1) = R(B(s)p_2)R(B(s)p_1) = B(s)R(p_2)R(p_1)B(-s) = R(p_2)R(p_1)$. We say that the points q_1, q_2 are obtained from p_1, p_2 by means of *bending*.

- The relation $R(p_2)R(p_1) = R(q_2)R(q_1)$ is called a *bending relation*.

It is easy to show that, modulo cancellations and orthogonal relations, the bending relations provide all relations with $n = 4$ (Corollary 3.3).

Dealing with a general relation $R(p_n) \dots R(p_2)R(p_1) = 1$, we can bend p_{j-1}, p_j , then bend p_{k-1}, p_k , etc. (the indices are modulo n). At some moment, we may arrive at $p_{j-1} = p_j$ or $\langle p_{j-1}, p_j \rangle = 0$ and cancel $R(p_j)R(p_{j-1})$ or apply an orthogonal relation, thus diminishing n . Also, with the use of cancellation and orthogonal relations, we can simultaneously alter the signs of p_{j-1}, p_j if the projective line $L(p_{j-1}, p_j)$ is hyperbolic. We expect that there is a finite number of *basic* relations that generate all the relations in the described way.

Denote by H_n the group generated by $r_1, r_2, \dots, r_n$ with the defining relations $r_j^2 = 1$, $j = 1, \dots, n$, and $r_n \dots r_2 r_1 = 1$. Every relation $R(p_n) \dots R(p_2)R(p_1) = 1$ gives rise to a representation $H_n \rightarrow \text{PU}V$. The most interesting are the faithful and discrete ones.

1.1. Conjecture. There exists a finite number of basic relations, each one induces a faithful and discrete representation that remains faithful and discrete after a finite number of bendings.

For such a representation, the projective line $L(p_{j-1}, p_j)$ cannot be spherical as $R(p_j)R(p_{j-1})$ would otherwise be elliptic. Nor does it rate to be euclidean since bending p_j, p_{j+1} usually leads to a spherical $L(p_{j-1}, p_j)$. Unfortunately, even if at the beginning all projective lines $L(p_{j-1}, p_j)$ are hyperbolic, some of them can become spherical after a few bendings (see Section 4.3). This will not happen if at most one of the p_j 's is positive.

As a first step, we study the representations

$$\varrho : H_5 \rightarrow \text{PU}V, \quad \varrho : r_j \mapsto R(p_j), \quad p_j \in \mathbb{P}_{\mathbb{C}}V \setminus SV$$

such that at most one of the p_j 's is positive. For brevity, we call them *pentagons*. Lifting a pentagon to $\text{SU}V$, we obtain the relation

$$R(p_5)R(p_4)R(p_3)R(p_2)R(p_1) = \delta,$$

where $\delta^3 = 1$. We show that, modulo conjugation, two pentagons with the same δ are connected by means of a finite number of bendings (see Subsection 5.4). In other words, each cubic root of unity δ provides a connected component of the space of pentagons. If $\delta = 1$, the pentagons are $\mathbb{R}$ -fuchsian and the p_j 's are negative. In this case, all pentagons are faithful and discrete [ABG]. Therefore, while deforming such a pentagon, there is always some real plane stable under the action of ϱH_5 and ϱ remains faithful and discrete. An example of a faithful and discrete pentagon with $\delta \neq 1$ is constructed in [Ana].

1.2. Conjecture. Every pentagon is faithful and discrete.

We believe that the situation with pentagons is quite similar to that in Toledo's rigidity theorem (see [Tol] and [Gol1]). This analogy would be deeper if one could find a holomorphic section of the example in [Ana]. In this case, the corresponding holomorphic disc in $\mathbb{H}_{\mathbb{C}}^2$ would be stable under the action of H_5 .

We can describe bendings in the following more general terms. Let $p \in \mathbb{P}_{\mathbb{C}}V \setminus SV$. The bundle $\text{PU}V \rightarrow \text{PU}V/\text{Stab } p$, where $\text{PU}V/\text{Stab } p = \text{BV}$ if p is negative and $\text{PU}V/\text{Stab } p = \text{EV}$ if p is positive, is endowed with the (Cartan) connection whose horizontal subspaces are orthogonal to cosets with respect to the Killing form. In other words, any piecewise smooth path $c : [a, b] \rightarrow \mathbb{P}_{\mathbb{C}}V \setminus SV$ uniquely lifts to a path $F_c : [a, b] \rightarrow \text{PU}V$ such that $F_c(s)c(a) = c(s)$ and $F_c(a) = 1$. We call F_c the *isometry following c* . The concept of path-following isometry can be illustrated with a snail that walks along the path c . During the walk, the ambient geometry may change from the snail viewpoint and the snail shell may not suit anymore a new geometry. Nevertheless, the snail takes no risk to break its shell even if the shell is as large as the whole space $\mathbb{H}_{\mathbb{C}}^2 = \text{BV}$ or EV (see Proposition 2.4 for the list of basic properties of F_c). It is worthwhile mentioning that this concept is applicable to any classic geometry (see [AGr]). The isometry following a (suitably parameterized) geodesic $G\langle p_1, p_2 \rangle$ is exactly the bending $B(s)$ involving the points p_1, p_2 (see Proposition 2.6).

The main result of this paper is Theorem 4.2.2. Briefly speaking, it claims that a generic (i.e., regular in the sense of Definition 3.6) isometry $F \in \text{SU}V$ is a product of 3 reflections, $F = R(p_3)R(p_2)R(p_1)$, that two such decompositions are connected by means of finitely many bendings involving $p_1, p_2/p_2, p_3$, and that geometrically equal decompositions differ by an isometry centralizing F . (The case when p_1, p_2, p_3 are in a same real plane is simpler and slightly different.) A certain elementary exposition [ABG] of a part of the classic Teichmüller space theory is intended to be applicable to the complex hyperbolic plane. In this sense, Theorem 4.2.2 is a first move towards studying the Teichmüller space $\mathcal{TH}_n$ of the hyperelliptic group H_n for the complex hyperbolic plane. Following the line of the proof of Theorem 4.2.2, it seems possible to generalize it for > 3 reflections. Moreover, there may exist a variant of Theorem 4.2.2 valid for an arbitrary classic geometry [AGr].

In subsequent articles, we plan to study the Teichmüller space $\mathcal{TH}_n$ in detail. In particular, the path-following isometry is intended to provide Toledo-like discrete invariants of discrete representations and to distinguish some of the examples constructed in [AGG] that have the same topology and the same Toledo invariant.

2. Path-following isometries

In this section, we introduce path-following isometries and describe bendings, i.e., isometries following geodesics.

We use the conventions and notation from [AGr] and [AGG]. Depending on the context, the elements in V may denote points in $\mathbb{P}_{\mathbb{C}}V$. For the sign of a point $p \in \mathbb{P}_{\mathbb{C}}V$, we use the notation $\sigma p \in \{-1, 0, 1\}$. In what follows, $\dot{c}(s)$ stands for the derivative of $c(s)$ with respect to s . We hope that the same notation for tangent vectors to curves in V and in $\mathbb{P}_{\mathbb{C}}V$ will produce no confusion since these tangent vectors ‘live’ in totally different places.

2.1. Remark. Let $c : [a, b] \rightarrow \mathbb{P}_{\mathbb{C}}V \setminus SV$ be a piecewise smooth path. Then there exists a piecewise smooth lift $c_0 : [a, b] \rightarrow V$ such that $\langle c_0(s), c_0(s) \rangle = \sigma \in \{-1, +1\}$ and $\langle c_0(s), \dot{c}_0(s) \rangle = 0$ for all $s \in [a, b]$. Such a lift is unique modulo multiplication by a unitary constant $u \in \mathbb{C}$, $|u| = 1$.

Proof. It is easy to find a piecewise smooth lift with $\langle c_0(s), c_0(s) \rangle = \sigma \in \{-1, +1\}$. It follows that $\langle c_0(s), \dot{c}_0(s) \rangle = if(s)$ for all $s \in [a, b]$, where $f : [a, b] \rightarrow \mathbb{R}$ is piecewise continuous. We can find a piecewise smooth function $\varphi : [a, b] \rightarrow \mathbb{R}$ such that $\dot{\varphi}(s)\sigma = f(s)$ for all $s \in [a, b]$. Now

$$\begin{aligned} \left\langle e^{i\varphi(s)}c_0(s), (e^{i\varphi(s)}c_0(s))' \right\rangle &= \langle c_0(s), \dot{c}_0(s) \rangle + \langle e^{i\varphi(s)}c_0(s), e^{i\varphi(s)}i\dot{\varphi}(s)c_0(s) \rangle = \\ &= i\left(f(s) - \dot{\varphi}(s)\langle c_0(s), c_0(s) \rangle\right) = 0. \end{aligned}$$

For another lift $g(s)c_0(s)$, $g : [a, b] \rightarrow \mathbb{C}$, it follows from $\langle g(s)c_0(s), g(s)c_0(s) \rangle = \sigma$ that $|g(s)| = 1$ and, from $\langle g(s)c_0(s), (g(s)c_0(s))' \rangle = 0$, we deduce $g(s)\dot{g}(s)\sigma = 0$ and, hence, $\dot{g}(s) = 0$ ■

The lift c_0 in Remark 2.1 is said to be *normalized*.

Note that a normalized lift of a closed path can be nonclosed. For instance, for orthonormal positive $p, p' \in V$, put $c_0(s) := p \cos s + p' \sin s$, $s \in [0, \pi]$ (see Proposition 2.6). Clearly, $\langle c_0(s), c_0(s) \rangle = 1$, $\dot{c}_0(s) = -p \sin s + p' \cos s$, and $\langle c_0(s), \dot{c}_0(s) \rangle = 0$. So, c is a closed path, whereas $c_0(\pi) = -c_0(0)$.

In fact, we obtain a $U(1)$ -connection in the tautological line bundle over $\mathbb{P}_{\mathbb{C}}V \setminus SV$.

2.2. Remark. Let $c : [a, b] \rightarrow \mathbb{P}_{\mathbb{C}}V \setminus SV$ be a piecewise smooth path and let $c_0 : [a, b] \rightarrow V$ be a normalized lift of c . Then $\dot{c}(t) = \langle -, c_0(t) \rangle \sigma \dot{c}_0(t)$ by [AGG, Lemma 4.1.4] ■

The group $SU V$ is given by the ‘equations’ $\det X = 1$ and $\langle Xv_1, Xv_2 \rangle = \langle v_1, v_2 \rangle$, $v_1, v_2 \in V$. As the Lie algebra $\mathfrak{su} V$ is the tangent space $T_1 SU V$, it is given inside $\text{Lin}_{\mathbb{C}}(V, V)$ by the ‘equations’ $\text{tr} Y = 0$ and $\langle Yv_1, v_2 \rangle + \langle v_1, Yv_2 \rangle = 0$, $v_1, v_2 \in V$, which can be also written as $\text{tr} Y = 0$ and $Y + Y^* = 0$. Identifying $\text{Lin}_{\mathbb{C}}(V, V) \simeq V^* \otimes_{\mathbb{C}} V$, we have $Y^* = \langle -, v \rangle p$ for $Y := \langle -, p \rangle v$. So, $\text{tr} Y = \text{tr} Y^*$ if $\langle v, p \rangle \in \mathbb{R}$. Clearly, $T_F SU V = F \cdot \mathfrak{su} V = \mathfrak{su} V \cdot F$ for every $F \in SU V$.

2.3. Definition. For $p \in \mathbb{P}_{\mathbb{C}}V \setminus SV$ and $t \in T_p \mathbb{P}_{\mathbb{C}}V \subset \text{Lin}_{\mathbb{C}}(V, V)$, denote $\hat{t} := t - t^* \in \mathfrak{su} V$. Given a piecewise smooth path $c : [a, b] \rightarrow \mathbb{P}_{\mathbb{C}}V \setminus SV$ and some $F_0 \in SU V$, we have a piecewise continuous path $\hat{c} : [a, b] \rightarrow \mathfrak{su} V$. The Cauchy problem $\dot{F} = \hat{c} \cdot F$, $F(a) = F_0$, has a unique piecewise smooth solution $F : [a, b] \rightarrow SU V$. We call F the *isometry following the path c and beginning with F_0* . Denote by F_c the isometry that follows c and begins with 1.

Denote by $c \cup c' : [a, d] \rightarrow X$ the concatenation of paths $c : [a, b] \rightarrow X$ and $c' : [b, d] \rightarrow X$ such that $c(b) = c'(b)$.

2.4. Proposition. Let $c : [a, b] \rightarrow \mathbb{P}_{\mathbb{C}}V \setminus SV$ and $c' : [b, d] \rightarrow \mathbb{P}_{\mathbb{C}}V \setminus SV$ be piecewise smooth paths such that $c(b) = c'(b)$, let $F_0 \in SU V$, and let $\varphi, \psi : [a', b'] \rightarrow [a, b]$ be piecewise smooth reparameterizations of c such that $\varphi(a') = a$, $\varphi(b') = b$ and $\psi(a') = b$, $\psi(b') = a$. Denote by F the isometry that follows c and begins with F_0 . Then

- $F(s) = F_c(s)F_0$ for all $s \in [a, b]$,
- $F_{c \cup c'} = F_c \cup F_{c'}F_c(b)$,
- $F_c \circ \varphi = F_{c \circ \varphi}$,
- $F_c \circ \psi = F_{c \circ \psi}F_c(b)$,
- $F_c(s)(c_0(a)) = c_0(s)$ for all $s \in [a, b]$, where $c_0 : [a, b] \rightarrow V$ stands for a normalized lift of c .

Proof. We will prove only the last two statements. For $\alpha := c \circ \psi$, we have $\dot{\alpha}(s) = \dot{c}(\psi(s))\dot{\psi}(s)$ and $\hat{\alpha}(s) = \hat{c}(\psi(s))\dot{\psi}(s)$. Since $(F_c \circ \psi)'(s) = \dot{F}_c(\psi(s))\dot{\psi}(s) = \hat{c}(\psi(s))F_c(\psi(s))\dot{\psi}(s) = \hat{\alpha}(s)(F_c \circ \psi)(s)$ and $(F_c \circ \psi)(a') = F_c(b)$, the fact follows.

For $\beta(s) := F_c(s)(c_0(a))$, we have

$$\begin{aligned} \dot{\beta}(s) &= \dot{F}_c(s)(c_0(a)) = \hat{c}(s)F_c(s)(c_0(a)) = \left(\dot{c}(s) - (\dot{c}(s))^* \right) \beta(s) = \\ &= \left(\langle -, c_0(s) \rangle \sigma \dot{c}_0(s) - \langle -, \dot{c}_0(s) \rangle \sigma c_0(s) \right) \beta(s) = \langle \beta(s), c_0(s) \rangle \sigma \dot{c}_0(s) - \langle \beta(s), \dot{c}_0(s) \rangle \sigma c_0(s) \end{aligned}$$

by Remark 2.2. It remains to observe that both $\beta(s)$ and $c_0(s)$ serve as solutions for the Cauchy problem $\dot{y}(s) = \langle y(s), c_0(s) \rangle \sigma \dot{c}_0(s) - \langle y(s), \dot{c}_0(s) \rangle \sigma c_0(s)$, $y(a) = c_0(a)$, for an unknown $y : [a, b] \rightarrow V$ ■

2.5. Lemma. Let $p \in \mathbb{P}_{\mathbb{C}}V \setminus SV$, let $c : [a, b] \rightarrow \mathbb{P}_{\mathbb{C}}V \setminus SV$ be a piecewise smooth path, and let $c(a) = F_0 p$ with $F_0 \in SU V$. Then the isometry $F : [a, b] \rightarrow SU V$ that follows c and begins with F_0 is a

horizontal lift of c with respect to the connection in the bundle $\mathrm{SU} V \rightarrow \mathrm{SU} V / \mathrm{Stab} p$ whose horizontal spaces are the tangent subspaces orthogonal to cosets with respect to the Killing form.

Proof. For a nonisotropic q , the subspace $\widehat{T_q \mathbb{P}_{\mathbb{C}} V} \subset \mathrm{su} V$ coincides with $(T_1 \mathrm{Stab} q)^\perp$. Indeed, let $l \in T_1 \mathrm{Stab} q$ and let $t \in T_q \mathbb{P}_{\mathbb{C}} V$, that is, $t = \langle -, q \rangle v$, $\langle v, q \rangle = 0$, $l \in \mathrm{su} V$, and $lq = 0$. Then $\mathrm{tr}(\widehat{lt}) = \langle lv, q \rangle = -\langle v, lq \rangle = 0$. It remains to compare the dimensions of $\widehat{T_q \mathbb{P}_{\mathbb{C}} V}$ and $(T_1 \mathrm{Stab} q)^\perp$.

In particular, $\dot{F}(s) \in (T_1 \mathrm{Stab} c(s))^\perp \cdot F(s)$ for any $s \in [a, b]$. By Proposition 2.4, $F(s) = F_c(s)F_0$ and $F(s)p = c(s)$. Hence, $(\mathrm{Stab} c(s))F(s) = F(s) \mathrm{Stab} p$ and $\dot{F}(s) \in F(s) \cdot (T_1 \mathrm{Stab} p)^\perp$. ■

We will describe isometries following geodesics. Due to Proposition 2.4, we should not worry much about the way of parameterizing geodesics. So, we use the most convenient parameter. The *bending* $B(s)$ involving p_1, p_2 is in fact the isometry following the geodesic $G \langle p_1, p_2 \rangle$ taken with a certain parameterization proportional to the natural one. It is explicitly defined in the proof of the following proposition.

2.6. Proposition. *Let $p_1, p_2 \in \mathbb{P}_{\mathbb{C}} V \setminus SV$ be distinct and nonorthogonal. All solutions $x_1, x_2 \in \mathbb{P}_{\mathbb{C}} V \setminus SV$ of the equation $R(x_2)R(x_1) = R(p_2)R(p_1)$ can be described as follows.*

There exists a one-parameter subgroup $B : (\mathbb{R}, +) \rightarrow \mathrm{SU} V$ called bending involving p_1, p_2 such that $B(s)$ commutes with $R(p_2)R(p_1)$ and $B(s)G = G$ for all $s \in \mathbb{R}$, where $G := G \langle p_1, p_2 \rangle$ stands for the geodesic joining p_1, p_2 . If p_1, p_2 have the same sign, then $B(1)p_1 = p_2$. If p_1, p_2 have different signs, the point $B(1)p_1 \in G$ is orthogonal to p_2 .

For an arbitrary $s \in \mathbb{R}$, we define $q_j := B(s)p_j$, $j = 1, 2$. Then $R(q_2)R(q_1) = R(p_2)R(p_1)$. If G is noneuclidean, we denote by $q'_j \in G$ the point orthogonal to $q_j \in G$, $j = 1, 2$. Then $R(q'_2)R(q'_1) = R(p_2)R(p_1)$.

Proof. Denote by p the polar point to $L := L(p_1, p_2)$. Note that the points x_1, x_2 subject to $R(x_2)R(x_1) = R(p_2)R(p_1)$ cannot be equal or orthogonal by Remark 3.2. If L is noneuclidean, the equality $R(q_2)R(q_1) = R(p_2)R(p_1)$ implies the equality $R(q'_2)R(q'_1) = R(p_2)R(p_1)$. Indeed, using the orthogonal relations (see Remark 3.2), we obtain $R(p_2)R(p_1) = R(q_2)R(q_1) = R(q_2)R(p)R(p)R(q_1) = R(q'_2)R(q'_1)$.

• Suppose that G is hyperbolic. Choose representatives of the vertices $v_1, v_2 \in G \cap SV$ such that the Gram matrix of v_1, v_2, p equals $\begin{bmatrix} 0 & \frac{1}{2} & 0 \\ \frac{1}{2} & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$, $p_1 = v_1 + \sigma_1 v_2$, and $p_2 = e^{-a}v_1 + \sigma_2 e^a v_2$, $0 \neq a \in \mathbb{R}$, where $\sigma_j := \sigma p_j$, $j = 1, 2$. The positive/negative points in G have the form $c_\pm(s) := e^{-as}v_1 \pm e^{as}v_2$, $s \in \mathbb{R}$. It is easy to see that $\langle c_\pm(s), c_\pm(s) \rangle = \pm 1$ and $\langle c_\pm(s), \dot{c}_\pm(s) \rangle = 0$. So, the lifts are normalized. By Remark 2.2, $\dot{c}_\pm(s) = \pm \langle -, c_\pm(s) \rangle \dot{c}_\pm(s)$. It follows that $\widehat{c}(s) := \widehat{c}_+(s) = \widehat{c}_-(s) = \begin{bmatrix} -a & 0 & 0 \\ 0 & a & 0 \\ 0 & 0 & 0 \end{bmatrix}$. Obviously,

$$F(s) := F_{c_+}(s) = F_{c_-}(s) = \begin{bmatrix} e^{-as} & 0 & 0 \\ 0 & e^{as} & 0 \\ 0 & 0 & 1 \end{bmatrix} \in \mathrm{SU}(2, 1) \text{ meets the conditions } \dot{F}(s) = \widehat{c}(s)F(s) \text{ and } F(0) = 1.$$

We define $B(t) := F(t)$ and call the one-parametric subgroup $B : (\mathbb{R}, +) \rightarrow \mathrm{SU} V$ the *bending involving* p_1, p_2 . Note that $B(1)p_1 = p_2$ if $\sigma_1 = \sigma_2$ and $B(1)p_1 \in L$ is orthogonal to $p_2 \in L$ if $\sigma_1 \neq \sigma_2$.

It is easy to verify that $B(s')c_\pm(s) = c_\pm(s + s')$ and $R(c_\pm(s)) = \begin{bmatrix} 0 & \pm e^{-2as} & 0 \\ \pm e^{2as} & 0 & 0 \\ 0 & 0 & -1 \end{bmatrix}$ for all $s, s' \in \mathbb{R}$.

$$\text{So, } R(c_{\sigma_2}(s_2))R(c_{\sigma_1}(s_1)) = \begin{bmatrix} \sigma_1 \sigma_2 e^{-2a(s_2 - s_1)} & 0 & 0 \\ 0 & \sigma_1 \sigma_2 e^{2a(s_2 - s_1)} & 0 \\ 0 & 0 & 1 \end{bmatrix} \text{ and } R(q_2)R(q_1) = R(p_2)R(p_1).$$

For the converse, it suffices to observe that the points $x_1, x_2 \in \mathbb{P}_{\mathbb{C}} V \setminus SV$ meeting the equation $R(x_2)R(x_1) = R(p_2)R(p_1)$ have to belong to the geodesic G because G is determined by the fixed points v_1, v_2, p of $R(p_2)R(p_1)$.

• Suppose that G is spherical. Let $p'_1 \in G$ be such that p_1, p'_1 are orthonormal. For every fixed $0 \neq a \in \mathbb{R}$, the points in G have the form $c(s) := p_1 \cos as + p'_1 \sin as$, $s \in \mathbb{R}$. It is easy to see that

this formula provides a normalized lift of c . Choose $a \in (0, \frac{\pi}{2})$ and representatives of p_1, p'_1, p_2 such that p_1, p'_1 are orthonormal and $c(1) = p_2$. By Remark 2.2, $\dot{c}(s) = \langle -, c(s) \rangle \dot{c}(t)$. In the orthonormal basis p_1, p'_1, p , we have $\hat{c}(s) = \begin{bmatrix} 0 & -a & 0 \\ a & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$. So, $F_c(s) = \begin{bmatrix} \cos as & -\sin as & 0 \\ \sin as & \cos as & 0 \\ 0 & 0 & 1 \end{bmatrix} \in \text{SU}(2, 1)$ meets the conditions $\dot{F}_c(s) = \hat{c}(s)F_c(s)$ and $F_c(0) = 1$.

We define $B(s) := F_c(s)$ and call the one-parametric subgroup $B : (\mathbb{R}, +) \rightarrow \text{SU} V$ the *bending involving* p_1, p_2 . Obviously, $B(1)p_1 = p_2$.

It is easy to verify that $B(s')c(s) = c(s + s')$ and $R(c(s)) = \begin{bmatrix} \cos 2as & \sin 2as & 0 \\ \sin 2as & -\cos 2as & 0 \\ 0 & 0 & -1 \end{bmatrix}$. This implies that $R(c(s_2))R(c(s_1)) = \begin{bmatrix} \cos 2a(s_2 - s_1) & -\sin 2a(s_2 - s_1) & 0 \\ \sin 2a(s_2 - s_1) & \cos 2a(s_2 - s_1) & 0 \\ 0 & 0 & 1 \end{bmatrix}$ and $R(q_2)R(q_1) = R(p_2)R(p_1)$.

It remains to observe that the fixed points $p_1 + ip'_1, p_1 - ip'_1, p$ of $R(p_2)R(p_1)$ determine² the geodesic G since $G = \{g \in L \mid \text{ta}(g, p + ip') = \text{ta}(g, p - ip')\}$.

• Suppose that G is euclidean. Choose $b \in BV \cap \mathbb{P}_{\mathbb{C}} p_1^\perp$ and representatives of b, p_1, p, p_2 such that the Gram matrix of b, p_1, p equals $\begin{bmatrix} -1 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}$ and $p_2 = p_1 + ap$ for some $a > 0$. Every point in G but p has the form $c_0(s) := p_1 + asp$, $s \in \mathbb{R}$. It is easy to see that another choice of b and the representatives does not change the parameterization $c(s)$ of G (however, it may change a). Since $\langle c_0(s), c_0(s) \rangle = 1$ and $\langle c_0(s), \dot{c}_0(s) \rangle = 0$, the lift c_0 is normalized. By Remark 2.2, $\dot{c}(s) = \langle -, c_0(s) \rangle \dot{c}_0(s) = \langle -, c_0(s) \rangle ap$. It follows that $\hat{c}_0(s) = \begin{bmatrix} 0 & 0 & 0 \\ -a & 0 & 0 \\ 0 & a & 0 \end{bmatrix}$ in the basis b, p_1, p . Therefore, $F_c(s) = \begin{bmatrix} 1 & 0 & 0 \\ -as & 1 & 0 \\ -a^2s^2/2 & as & 1 \end{bmatrix} \in \text{SU}(2, 1)$ meets the conditions $\dot{F}_c(s) = \hat{c}(s)F_c(s)$ and $F_c(0) = 1$.

We define $B(s) := F_c(s)$ and call the one-parametric subgroup $B : (\mathbb{R}, +) \rightarrow \text{SU}(2, 1)$ the *bending involving* p_1, p_2 . Obviously, $B(1)p_1 = p_2$.

A straightforward verification shows that $B(s')c_0(s) = c_0(s + s')$ and $R(c_0(s)) = \begin{bmatrix} -1 & 0 & 0 \\ 2as & 1 & 0 \\ 2a^2s^2 & 2as & -1 \end{bmatrix}$. Hence, $R(c_0(s_2))R(c_0(s_1)) = \begin{bmatrix} 1 & 0 & 0 \\ -2a(s_2 - s_1) & 1 & 0 \\ -2a^2(s_2 - s_1)^2 & 2a(s_2 - s_1) & 1 \end{bmatrix}$. So, $R(q_2)R(q_1) = R(p_2)R(p_1)$ and these are unique solutions $x_1, x_2 \in G$ of the equation $R(x_2)R(x_1) = R(p_2)R(p_1)$.

As p is the only fixed point of $R(p_2)R(p_1)$, the points x_1, x_2 subject to the equation $R(x_2)R(x_1) = R(p_2)R(p_1)$ should belong to L . The geodesic G is stable under the action of $R(p_2)R(p_1)$. So is the geodesic $G \setminus \{x_1, x_2\} \subset L$. Writing down the x_j 's in the form $x_j = p_1 + c_j p$, $c_j \in \mathbb{C}$, we conclude that $p_1 + (2a + c_j)p \in G \setminus \{p_1 + c_1 p, p_1 + c_2 p\}$ and derive from this that $c_1, c_2 \in \mathbb{R}$ ■

It is not difficult to show that the slice identification introduced in [AGG] and the meridional transport mentioned in [AGr] are induced by suitable bendings. Also, one can derive the formulae for parallel transport along geodesics, given in [AGr], from the above description of bendings.

3. Bendings and reflections

In this section, we study how an arbitrary relation between reflections can be modified with the help of short relations ($n \leq 4$). Also, we show that almost all isometries in $\text{SU} V$ are products of 3 reflections.

Let $p \in \mathbb{P}_{\mathbb{C}} V \setminus SV$. The *reflection* $R(p) : V \rightarrow V$ in p is defined as

$$R(p) : x \mapsto 2 \frac{\langle x, p \rangle}{\langle p, p \rangle} p - x.$$

It is easy to see that $R(p) \in \text{SU} V$ and $R(p)R(p) = 1$.

²I am indebted to Carlos Henrique Grossi Ferreira for pointing out this nice and easy fact.

Given pairwise orthogonal $p_1, p_2, p_3 \in \mathbb{P}_{\mathbb{C}}V \setminus SV$, it is immediate that the $R(p_i)$'s commute and $R(p_3)R(p_2)R(p_1) = 1$.

Denote

$$\text{ta}(p_1, p_2) := \frac{g_{12}g_{21}}{g_{11}g_{22}}, \quad \alpha(p_1, p_2, p_3) := \text{Im} \frac{g_{12}g_{23}g_{31}}{g_{11}g_{22}g_{33}}, \quad \beta(p_1, p_2, p_3) := \frac{\det[g_{ji}]}{g_{11}g_{22}g_{33}}, \quad \tau(p_1, p_2, p_3) := \text{Re} \frac{g_{13}g_{22}}{g_{12}g_{23}},$$

where $[g_{jk}]$ stands for the Gram matrix of nonisotropic $p_1, p_2, p_3 \in V$, i.e., $g_{jk} := \langle p_j, p_k \rangle$.

3.1. Lemma (compare with [Pra]). *Let $[g_{jk}]$ be the Gram matrix of $p_1, \dots, p_n \in \mathbb{P}_{\mathbb{C}}V \setminus SV$. Then*

$$\text{tr}(R(p_n) \dots R(p_1)) = (-1)^n \left(3 - 2n + \sum_{\substack{1 \leq i_1 < \dots < i_t \leq n \\ 2 \leq t \leq n}} (-2)^t \frac{g_{i_1 i_2} g_{i_2 i_3} \dots g_{i_{t-1} i_t} g_{i_t i_1}}{g_{i_1 i_1} g_{i_2 i_2} \dots g_{i_{t-1} i_{t-1}} g_{i_t i_t}} \right).$$

In particular,

$$\text{tr}(R(p_1)) = -1, \quad \text{tr}(R(p_2)R(p_1)) = 4\text{ta}(p_1, p_2) - 1,$$

$$\text{tr}(R(p_3)R(p_2)R(p_1)) = 8i\alpha(p_1, p_2, p_3) + 4\beta(p_1, p_2, p_3) - 1.$$

Proof. The facts that the trace of $x \mapsto \langle x, q \rangle p$ equals $\langle p, q \rangle$ and that $\text{tr} 1 = 3$ imply the general formula. The rest follows from

$$\det[g_{ji}] = g_{11}g_{22}g_{33} + 2\text{Re}(g_{12}g_{23}g_{31}) - g_{13}g_{31}g_{22} - g_{12}g_{21}g_{33} - g_{23}g_{32}g_{11} =$$

$$= g_{11}g_{22}g_{33} \left(1 + 2 \frac{\text{Re}(g_{12}g_{23}g_{31})}{g_{11}g_{22}g_{33}} - \text{ta}(p_1, p_2) - \text{ta}(p_2, p_3) - \text{ta}(p_3, p_1) \right),$$

$$\text{tr}(R(p_3)R(p_2)R(p_1)) = 3 - 4(\text{ta}(p_1, p_2) + \text{ta}(p_2, p_3) + \text{ta}(p_3, p_1)) + 8 \frac{g_{12}g_{23}g_{31}}{g_{11}g_{22}g_{33}} \blacksquare$$

The following well-known facts are straightforward or easily follow from Lemma 3.1 :

3.2. Remark. *Let $\delta \in \mathbb{C}$ be such that $\delta^3 = 1$ and let $p_1, p_2, p_3 \in \mathbb{P}_{\mathbb{C}}V \setminus SV$.*

- *If $F^2 = \delta$ for some $F \in \text{SU} V$, then either $F = \delta^2$ or $F = \delta^2 R(p)$ for a suitable $p \in \mathbb{P}_{\mathbb{C}}V \setminus SV$.*
- *If $R(p_2)R(p_1) = \delta$, then $\delta = 1$ and $p_1 = p_2$.*
- *If $R(p_3)R(p_2)R(p_1) = \delta$, then $\delta = 1$ and p_1, p_2, p_3 are pairwise orthogonal* ■

Suppose that $R(q_1)R(p_1) = \delta R(q_2)R(p_2)$ with $\delta^3 = 1$. Then $\delta = 1$. Indeed, $4\text{ta}(p_1, q_1) - 1 = \delta(4\text{ta}(p_2, q_2) - 1)$ by Lemma 3.1. If $\delta \neq 1$, then $4\text{ta}(p_j, q_j) = 1$ and the projective line $L(p_j, q_j)$ is spherical, $j = 1, 2$. Hence, the polar point b_j to $L(p_j, q_j)$ is negative. Being b_j the only negative fixed point of $R(q_j)R(p_j)$, we obtain $b_1 = b_2$. The other two fixed points of $R(q_1)R(p_1)$ and of $R(q_2)R(p_2)$ also coincide. For the reason used in the proof of Proposition 2.6, $G \wr p_1, q_1 \wr = G \wr p_2, q_2 \wr$. Applying a suitable bending involving p_2, q_2 , we can assume that $p_2 = p_1$. A contradiction.

So, from Remark 3.2 and Proposition 2.6, we obtain the following corollary.

3.3. Corollary. *All relations with $n \leq 4$ follow from cancellations, orthogonal relations, and bending relations* ■

3.4. Remark. *Given $p_1, p_2, p_3 \in \mathbb{P}_{\mathbb{C}}V \setminus SV$ such that p_1, p_2 are distinct nonorthogonal and $L := L(p_1, p_2)$ is not spherical, the projective line $L(p_2, p_3)$ becomes hyperbolic after a suitable bending involving p_1, p_2 except in the following cases:*

- *L is euclidean and $p_3 \in L$,*
- *L is hyperbolic and p_3 is the polar point to L .*

Proof. Denote by p the polar point to L . We can assume that $\langle p_2, p_2 \rangle = \langle p_3, p_3 \rangle = 1$.

Suppose that L is euclidean and that $p_3 \notin L$, i.e., $\langle p, p_3 \rangle \neq 0$. Every point in $G\langle p_1, p_2 \rangle \setminus \{p\}$ has the form $p_2 + sp$, $s \in \mathbb{R}$, after a suitable choice of a representative $p \in V$. Hence, $\text{ta}(p_2 + sp, p_3) = |\langle p_2, p_3 \rangle + s\langle p, p_3 \rangle|^2 \rightarrow \infty$ while $s \rightarrow \infty$ because $\langle p, p_3 \rangle \neq 0$.

Suppose that L is hyperbolic. Every positive point in $G\langle p_1, p_2 \rangle$ has the form $p(s) = e^{-s}v_1 + e^s v_2$, $s \in \mathbb{R}$, where v_1, v_2 stand for representatives of the vertices of $G\langle p_1, p_2 \rangle$ such that $\langle v_1, v_2 \rangle = \frac{1}{2}$. Therefore, $\text{ta}(p(s), p_3) = |e^{-s}\langle v_1, p_3 \rangle + e^s\langle v_2, p_3 \rangle|^2 \rightarrow \infty$ while $s \rightarrow \pm\infty$ unless $\langle v_1, p_3 \rangle = \langle v_2, p_3 \rangle = 0$ ■

3.5. Corollary. *Let $R(p_n) \dots R(p_1) = \delta$, where $p_1, \dots, p_n \in \mathbb{P}_{\mathbb{C}}V \setminus SV$ and $\delta^3 = 1$. Then, using cancellations, orthogonal relations, and bendings, we can either diminish n or reach a situation where at most one of the p_j 's is positive unless every projective line $L(p_{j-1}, p_j)$ is spherical (the indices are modulo n).*

Proof. First, we show how to gain a negative p_j . If some $L(p_{j-1}, p_j)$ is hyperbolic, this is easy since, using orthogonal relations, we can simultaneously alter the signs of p_{j-1}, p_j , if necessary. So, we assume $L(p_1, p_2)$ euclidean. By Remark 3.4, we assume that $p_3 \in L(p_1, p_2)$, that is, $L(p_1, p_2) = L(p_2, p_3)$. In this way, we arrive at a situation where the p_j 's belong to the same euclidean projective line L . It remains to apply elementary euclidean plane arguments. (See the end of [AGr, Subsection 3.1] for the description of geodesics in an euclidean projective line.) If the geodesics $G\langle p_1, p_2 \rangle$ and $G\langle p_3, p_4 \rangle$ intersect, i.e., if they have a common point q different from the polar point to L , then we obtain $p_3 = q$ after a suitable bending involving p_3, p_4 . Hence, $p_3 \in G\langle p_1, p_2 \rangle$ and some bending involving p_2, p_3 provides $p_2 = p_1$, i.e., a cancellation. If $p_5 \in G\langle p_3, p_4 \rangle$, then we can get $p_3 = p_4$ with a bending involving p_4, p_5 . If $p_5 \notin G\langle p_3, p_4 \rangle$, a 'small' bending involving p_4, p_5 makes $G\langle p_1, p_2 \rangle$ and $G\langle p_3, p_4 \rangle$ intersect.

Without loss of generality, we assume $p_1, \dots, p_{j-1}$ negative and p_j, p_n positive, $1 < j < n$. If p_{j+1} is the polar point to $L(p_{j-1}, p_j)$, we can diminish n by using an orthogonal relation. Otherwise, by Remark 3.4, $L(p_j, p_{j+1})$ becomes hyperbolic after a bending involving p_{j-1}, p_j . It remains to alter the signs of p_j, p_{j+1} , if necessary ■

3.6. Definition. Let $d = 2$. A triple of points $p_1, p_2, p_3 \in \mathbb{P}_{\mathbb{C}}V \setminus SV$ is said to be *regular* if p_1, p_2, p_3 are not in a same geodesic, p_2 is not orthogonal to p_j for $j = 1, 3$, and at most one of p_1, p_2, p_3 is positive. A regular triple p_1, p_2, p_3 is *strongly regular* when p_1, p_2, p_3 are not in a same projective line and p_1, p_2, p_3 are all negative if they belong to a same real plane. An isometry $F \in \text{SU } V$ is called *regular* if $\dim_{\mathbb{C}}\{v \in V \mid Fv = cv\} \leq 1$ for any $c \in \mathbb{C}$.

3.7. Lemma. *Let p_1, p_2, p_3 be a regular triple. Then the isometry $F := R(p_3)R(p_2)R(p_1)$ is regular.*

Proof. Suppose that $\dim_{\mathbb{C}}\{v \in V \mid Fv = cv\} \geq 2$. Then $|c| = 1$. Note that $c \neq -1$. Otherwise, $F = R(p)$ is a reflection and p, p_1, p_2, p_3 are in a same geodesic by Proposition 2.6, contradicting the assumption that the triple is regular. We can pick $0 \neq v \in V$ such that $\langle v, p_1 \rangle = 0$ and $Fv = cv$. This means that $R(p_1)v = -v$ and, hence, $R(p_2)v + cR(p_3)v = 0$. In other words, assuming that $\sigma_j := \langle p_j, p_j \rangle \in \{-1, 1\}$, we obtain $2\sigma_2\langle v, p_2 \rangle p_2 + 2\sigma_3 c\langle v, p_3 \rangle p_3 = (1+c)v$. Since $1+c \neq 0$ and p_2, p_3 are $\mathbb{C}$ -linearly independent, we conclude that $v \in \mathbb{C}p_2 + \mathbb{C}p_3$. If v is proportional to one of p_2, p_3 , then $\langle p_2, p_3 \rangle = 0$, which is impossible for a regular triple. Hence, we can assume that $v = zp_2 + p_3$, for some $0 \neq z \in \mathbb{C}$. By a straightforward calculation, we obtain $\langle p_2, p_3 \rangle = \frac{\sigma_2(\bar{c}-1)\bar{z}}{2} = \frac{\sigma_3(1-c)}{2cz}$. Therefore, $|z| = 1$ and $\sigma_2 = \sigma_3$, implying $\sigma_2 = \sigma_3 = -1$ and $|\langle p_2, p_3 \rangle| < 1$. It follows that $\text{ta}(p_2, p_3) < 1$, a contradiction ■

3.8. Proposition. *An isometry $F \in \text{SU } V$ admits the form $F = R(p_3)R(p_2)R(p_1)$, where p_1, p_2, p_3 is a regular triple, iff F is regular and $\text{tr } F \neq -1$.*

Proof. If $\text{tr}(R(p_3)R(p_2)R(p_1)) = -1$, then $\alpha(p_1, p_2, p_3) = \beta(p_1, p_2, p_3) = 0$ by Lemma 3.1. Hence, p_1, p_2, p_3 are in a same geodesic.

Conversely, given a regular $F \in \text{SU } V$ such that $\text{tr } F \neq -1$, write $\text{tr } F = 8i\alpha + 4\beta - 1$ and $\beta = \mp|\beta|$, where $\alpha, \beta \in \mathbb{R}$. For sufficiently big $g_{12}, g_{23} > 1$, we find $t \in \mathbb{R}$ satisfying the equation

$$g_{12}^2 g_{23}^2 (t-1)^2 = (g_{12}^2 \pm 1)(g_{23}^2 \pm 1) - \beta - \frac{\alpha^2}{g_{12}^2 g_{23}^2}.$$

We have

$$\pm \det G_{\pm} = 1 + 2g_{12}^2 g_{23}^2 t \pm g_{12}^2 \pm g_{23}^2 - g_{12}^2 g_{23}^2 t^2 - \frac{\alpha^2}{g_{12}^2 g_{23}^2} = (g_{12}^2 \pm 1)(g_{23}^2 \pm 1) - g_{12}^2 g_{23}^2 (t-1)^2 - \frac{\alpha^2}{g_{12}^2 g_{23}^2} = \beta = \mp|\beta|,$$

where

$$G_{\pm} := \begin{bmatrix} -1 & g_{12} \pm g_{12} g_{23} t \mp \frac{\alpha i}{g_{12} g_{23}} \\ g_{12} & \pm 1 \\ \pm g_{12} g_{23} t \pm \frac{\alpha i}{g_{12} g_{23}} & g_{23} & -1 \end{bmatrix}.$$

By Sylvester's Criterion, there exist $p_1, p_2, p_3 \in V$ with the Gram matrix $G_{\pm}$ because $g_{12} > 1$ and $\det G_{\pm} \leq 0$. Note that p_1, p_2, p_3 are not in the same geodesic because otherwise $\alpha = \beta = 0$ and $\text{tr } F = -1$. It remains to apply Lemma 3.1, [Gol2, Theorem 6.2.4], and Lemma 3.7 ■

In the sequel, we will need the following well-known fact which can be obtained by straightforward calculations:

3.9. Remark. *Let $F \in \text{SU } V$ be a regular isometry, let $C_0(F) \leq \text{U } V$ stand for the centralizer of F in $\text{U } V$, and let $C(F) \leq \text{PU } V$ denote the image of $C_0(F)$. Then $C(F)$ is connected, commutative, and 2-dimensional. ■*

4. Composition of bendings

Let p_1, p_2, p_3 be a strongly regular triple (see Definition 3.6). In this section, we study compositions of two types of bendings: those involving p_1, p_2 and those involving p_2, p_3 .

These bendings preserve the regular isometry $F := R(p_3)R(p_2)R(p_1)$ and the numbers $\sigma_j := \sigma p_j$, $j = 1, 2, 3$. So, we fix the σ_j 's and $\alpha, \beta \in \mathbb{R}$ such that $\text{tr } F = 8i\alpha + 4\beta - 1$, $\alpha := \alpha(p_1, p_2, p_3)$, and $\beta := \beta(p_1, p_2, p_3)$ (see Lemma 3.1). By Definition 3.6, Lemma 3.1, Proposition 3.8, and Sylvester's Criterion, the only restrictions on the numbers $\sigma_1, \sigma_2, \sigma_3, \alpha, \beta$ are

- $\sigma_1 \sigma_2 \sigma_3 \beta < 0$, at most one of the σ_j 's is positive, and $\alpha = 0 \implies \sigma_1 = \sigma_2 = \sigma_3 = -1$.

We will frequently choose representatives of p_1, p_2, p_3 such that $g_{jj} = \sigma_j$ and $g_{12}, g_{23} > 0$, where $[g_{jk}]$ stands for the Gram matrix of p_1, p_2, p_3 . Such a Gram matrix is called *standard*. Note that the standard Gram matrix is uniquely determined by a strongly regular triple $p_1, p_2, p_3 \in \mathbb{P}_{\mathbb{C}} V$.

When $\alpha = 0$, the standard Gram matrix is real. In this case, the triple lies in (and spans) a real plane. Such a triple is said to be *real*.

4.1. Geometrical configurations of strongly regular triples. First, we will show that, geometrically, i.e., up to the action of $\text{PU } V$, the strongly regular triples p_1, p_2, p_3 with fixed $\sigma_1, \sigma_2, \sigma_3$ and the conjugacy class of $F := R(p_3)R(p_2)R(p_1)$ form a surface S . Then we describe the lines on S that correspond to the bendings involving $p_1, p_2/p_2, p_3$.

4.1.1. Lemma. *Geometrically, a strongly regular triple p_1, p_2, p_3 can be described as a point in the surface $S \subset \mathbb{R}^3(t, t_1, t_2)$ given by the equation*

$$(4.1.2) \quad (t_1 - 1)(t_2 - 1) = t_1 t_2 (t - 1)^2 + \frac{\alpha^2}{t_1 t_2} + \beta$$

and by the inequalities

$$(4.1.3) \quad \sigma_1 \sigma_2 t_1 > 0, \quad \sigma_1 \sigma_2 t_1 > \sigma_1 \sigma_2, \quad \sigma_2 \sigma_3 t_2 > 0, \quad \sigma_2 \sigma_3 t_2 > \sigma_2 \sigma_3,$$

where $t_1 := \text{ta}(p_1, p_2)$, $t_2 := \text{ta}(p_2, p_3)$, and $t := \tau(p_1, p_2, p_3) := \text{Re} \frac{g_{13}g_{22}}{g_{12}g_{23}}$.

Proof. By Sylvester's Criterion, a strongly regular triple p_1, p_2, p_3 is geometrically given by its Gram matrix $[g_{jk}]$. We assume it to be standard. Let $t_1 := \text{ta}(p_1, p_2)$, $t_2 := \text{ta}(p_2, p_3)$, and $t := \tau(p_1, p_2, p_3) := \text{Re} \frac{g_{13}g_{22}}{g_{12}g_{23}}$. It follows from $\alpha(p_1, p_2, p_3) = \alpha$ that $g_{31} = \frac{g_{12}g_{23}}{g_{22}}t + \frac{g_{11}g_{22}g_{33}}{g_{12}g_{23}}\alpha i$. Therefore,

$$\beta := \frac{\det[g_{jk}]}{g_{11}g_{22}g_{33}} = 1 + 2t_1t_2 - t_1 - t_2 - \frac{g_{12}^2g_{23}^2}{g_{11}g_{22}g_{33}}t^2 - \frac{g_{11}g_{22}^2g_{33}}{g_{12}^2g_{23}}\alpha^2 = 1 + 2t_1t_2 - t_1 - t_2 - t_1t_2t^2 - \frac{\alpha^2}{t_1t_2}$$

and we arrive at (4.1.2). The inequalities $\sigma_1\sigma_2t_1 > \sigma_1\sigma_2$ and $\sigma_2\sigma_3t_2 > \sigma_2\sigma_3$ follow from the fact that the projective lines $L(p_1, p_2)$ and $L(p_2, p_3)$ are hyperbolic.

Conversely, let $(t, t_1, t_2) \in S$. We put $g_{jj} := \sigma_j$ for $j = 1, 2, 3$, $g_{12} := g_{21} := \sqrt{\sigma_1\sigma_2t_1} > 0$, $g_{23} := g_{32} := \sqrt{\sigma_2\sigma_3t_2} > 0$, $g_{31} := \frac{g_{12}g_{23}}{g_{22}}t + \frac{g_{11}g_{22}g_{33}}{g_{12}g_{23}}\alpha i$, and $g_{13} := \bar{g}_{31}$. The equation (4.1.2) implies that $\det[g_{jk}] = \sigma_1\sigma_2\sigma_3\beta$. By Sylvester's Criterion, there are points $p_1, p_2, p_3 \in V$ with the Gram matrix $[g_{jk}]$ because at most one of the σ_j 's is positive, $\sigma_1\sigma_2t_1 > \sigma_1\sigma_2$, and $\sigma_1\sigma_2\sigma_3\beta < 0$ ■

Given $r_1, r_2 \in \mathbb{R}$ such that $\sigma_1\sigma_2r_1 > 0$, $\sigma_1\sigma_2r_1 > \sigma_1\sigma_2$, $\sigma_2\sigma_3r_2 > 0$, and $\sigma_2\sigma_3r_2 > \sigma_2\sigma_3$, denote by V_{r_1} , H_{r_2} , and C the lines given in S respectively by the equations $t_1 = r_1$, $t_1 = r_2$, and $t = 1$. We call V_{r_1}/H_{r_2} a *vertical/horizontal* line. Obviously, the intersection of a vertical line and a horizontal one is either empty or a point in C or consists of two points outside C .

4.1.4. Lemma. *Every vertical/horizontal line is non-empty, smooth, and connected. It intersects C in exactly 1 point and corresponds to the bendings involving the points $p_1, p_2/p_2, p_3$. The surface S is a topological plane fibred by vertical/horizontal lines. If a vertical and a horizontal lines intersect outside C , this intersection is transversal.*

Proof. Let $Q \subset \mathbb{R}^2(t_1, t_2)$ denote the quadrant given by the inequalities (4.1.3). The projection of the surface S into the plane $\mathbb{R}^2(t_1, t_2)$ coincides with the region $R \subset Q$ given in Q by the inequality $f(t_1, t_2) \geq 0$, where $f(t_1, t_2) := \sigma_1\sigma_3((t_1 - 1)(t_2 - 1) - \frac{\alpha^2}{t_1t_2} - \beta)$. The map $S \rightarrow R$ is a double covering ramified along C . The fact that vertical/horizontal lines of S are smooth follows straightforwardly.³ Such lines project into the intersections with R of the vertical/horizontal lines of the plane $\mathbb{R}^2(t_1, t_2)$.

Let $\sigma_2\sigma_3r_2 > 0$ and $\sigma_2\sigma_3r_2 > \sigma_2\sigma_3$. The function


$$f(\sigma_1\sigma_2x, r_2) = (x - \sigma_1\sigma_2)(\sigma_2\sigma_3r_2 - \sigma_2\sigma_3) - \frac{\alpha^2}{\sigma_2\sigma_3r_2x} + \sigma_2|\beta|$$

is increasing in $x > 0$ and $\lim_{x \rightarrow +\infty} f(\sigma_1\sigma_2x, r_2) = +\infty$. Also, it takes negative values for $x > \max(\sigma_1\sigma_2, 0)$.

Indeed, if $\sigma_1\sigma_2 = 1$, then $\lim_{x \rightarrow +1} f(\sigma_1\sigma_2x, r_2) = -\frac{\alpha^2}{\sigma_2\sigma_3r_2} - |\beta|$. If $\sigma_1\sigma_2 = -1$, then $\alpha \neq 0$ and

$\lim_{x \rightarrow +0} f(\sigma_1\sigma_2x, r_2) = -\infty$. We conclude that every horizontal line of Q intersects C in exactly 1 point.

Hence, every horizontal line of S is connected and intersects C in exactly 1 point. By symmetry, the same is true for vertical lines.



³For example, if the derivatives of $r_1t_2(t-1)^2 + \frac{\alpha^2}{r_1t_2} + \beta - (r_1-1)(t_2-1)$ with respect to t and t_2 equal respectively $r_2(t-1)^2 - \frac{\alpha^2}{r_1t_2^2} - (r_1-1)$ and $2r_1t_2(t-1)$. If both vanish, then $t = 1$ and $\frac{\alpha^2}{t_2^2} + r_1(r_1-1) = 0$. On the other hand, $r_1(r_1-1) = \sigma_1\sigma_2r_1(\sigma_1\sigma_2r_2 - \sigma_1\sigma_2) > 0$.

It remains to prove the claim concerning the bendings. By symmetry, we deal only with vertical lines. Obviously, when we bend p_1, p_2 , we keep t_1 constant, $t_1 = r_1$. Denote by v_1, v_2 the vertices of the geodesic $G\langle p_1, p_2 \rangle$ and choose representatives of p_1, p_2, p_3, v_1, v_2 such that $\langle p_3, p_3 \rangle = \sigma_3$, $\langle v_1, v_2 \rangle = \frac{1}{2}$, $p_1 = v_1 + \sigma_1 v_2$, and $p_2 = e^{-a} v_1 + \sigma_2 e^a v_2$ with $a > 0$ and $a \neq 1$. The formulae

$$p_1(s) := e^{-as} v_1 + \sigma_1 e^{as} v_2, \quad p_2(s) := e^{-a(s+1)} v_1 + \sigma_2 e^{a(s+1)} v_2, \quad s \in \mathbb{R},$$

describe the bendings involving p_1, p_2 . Denote $z_j := \langle v_j, p_3 \rangle$ and $c_j := |z_j|$ for $j = 1, 2$. Note that $z_j \neq 0$. Indeed, suppose that $z_2 = 0$. Then $\sigma_3 = 1$, hence, $\sigma_1 = \sigma_2 = -1$ and $\alpha \neq 0$. On the other hand, $\alpha = \sigma_1 \sigma_2 \sigma_3 \operatorname{Im} \left(\frac{1}{2} (\sigma_1 e^{-a} + \sigma_2 e^a) e^{-a(2s+1)} z_1 \bar{z}_1 \right) = 0$, a contradiction.

From

$$t(s) = \operatorname{Re} \frac{2(e^{-as} z_1 + \sigma_1 e^{as} z_2) \sigma_2}{(\sigma_1 e^{-a} + \sigma_2 e^a)(e^{-a(s+1)} z_1 + \sigma_2 e^{a(s+1)} z_2)} = \frac{2(\sigma_2 e^{-a(2s+1)} c_1^2 + (\sigma_1 \sigma_2 e^{-a} + e^a) \operatorname{Re}(z_1 \bar{z}_2) + \sigma_1 e^{a(2s+1)} c_2^2)}{(\sigma_1 e^{-a} + \sigma_2 e^a) |e^{-a(s+1)} z_1 + \sigma_2 e^{a(s+1)} z_2|^2},$$

we obtain

$$\lim_{s \rightarrow -\infty} t(s) + \lim_{s \rightarrow +\infty} t(s) = \frac{2\sigma_2}{(\sigma_1 e^{-a} + \sigma_2 e^a) e^{-a}} + \frac{2\sigma_1}{(\sigma_1 e^{-a} + \sigma_2 e^a) e^a} = 2.$$

So, $\lim_{s \rightarrow -\infty} t(s) - 1$ and $\lim_{s \rightarrow +\infty} t(s) - 1$ have opposite signs. On the other hand, the function

$$\sigma_2 \sigma_3 t_2(s) = |e^{-a(s+1)} z_1 + \sigma_2 e^{a(s+1)} z_2|^2 = e^{-2a(s+1)} c_1^2 + e^{2a(s+1)} c_2^2 + 2\sigma_2 \operatorname{Re}(z_1 \bar{z}_2)$$

in s possesses exactly one minimum and $\lim_{s \rightarrow \pm\infty} \sigma_2 \sigma_3 t_2(s) = +\infty$. Taking into account the above description of a vertical line and its projection into Q , this completes the proof ■

4.1.5. Corollary. *The bending involving $p_1, p_2/p_2, p_3$ keeps any strongly regular triple p_1, p_2, p_3 strongly regular. Geometrically, all strongly regular triples with fixed α , β , and σ_j 's are connected by means of such bendings ■*

4.2. Holonomy of strongly regular triples. It can easily happen that, after a few bendings involving $p_1, p_2/p_2, p_3$, we obtain from a strongly regular triple p_1, p_2, p_3 a different triple p'_1, p'_2, p'_3 which is geometrically the same. In order to study this phenomena, we introduce some formal settings.

Denote by E the manifold of all strongly regular triples $p_1, p_2, p_3 \in \mathbb{P}_{\mathbb{C}} V$ (the conjugacy class of $R(p_3)R(p_2)R(p_1) \in \operatorname{SU} V$ and the σ_j 's are fixed). To every such triple, we can associate its standard Gram matrix; this matrix determines a triple of representatives $p_1, p_2, p_3 \in V$, unique up to a unitary factor. We obtain the map $\pi : E \rightarrow S$. By Sylvester's Criterion, π is a principal $\operatorname{PU} V$ -bundle.

To every triple $p := (p_1, p_2, p_3) \in E$, we associate the isometry $F_p := R(p_3)R(p_2)R(p_1) \in \operatorname{SU} V$. The left action of $\operatorname{PU} V$ on E induces the conjugation at the level of F_p . Indeed, let $g \in \operatorname{PU} V$ be a representative of $g \in \operatorname{PU} V$. Then

$$F_{gp} = R(gp_3)R(gp_2)R(gp_1) = gR(p_3)R(p_2)R(p_1)g^{-1} = gF_p g^{-1}.$$

In particular, the triples p with fixed $F_p = F$ form a principal $C(F)$ -bundle $E_F \rightarrow S$.

By Lemma 4.1.4, every vertical/horizontal line in S lifts to the trajectory of the corresponding bending. This provides a connection over $S \setminus C$ because at any point $q \in S \setminus C$ the tangent vectors to the vertical and horizontal lines passing through q generate the tangent space $T_q S$. (However, at a point in C , the vertical and horizontal lines are tangent.) Denote by $H(q_0)$ the corresponding (connected) holonomy group at $q_0 \in S \setminus C$ and, for any $p \in \pi^{-1}(q_0)$, let

$$H(p) := \{g \in \operatorname{PU} V \mid gp = ph \text{ for some } h \in H(q_0)\} \leq \operatorname{PU} V$$

stand for the corresponding subgroup. (By convention, $H(q_0)$ acts on $\pi^{-1}(q_0)$ from the right and this action commutes with that of $\text{PU}V$.)

Consider a piecewise vertical-horizontal path $c : [a, b] \rightarrow S$ and its lift $\tilde{c} : [a, b] \rightarrow E$, $\pi \circ \tilde{c} = c$, that begins at an arbitrary $p \in \pi^{-1}(c(a))$, $\tilde{c}(a) = p$. By construction, the path $\tilde{c}$ lies entirely in some E_F . Denote by $\varphi_c : [a, b] \rightarrow \text{PGL}V$ the path given by $p\varphi_c(s) = \tilde{c}(s)$ in terms of representatives $p_1, p_2, p_3 \in V$ with standard Gram matrix such that $p = (p_1, p_2, p_3)$, where $(p_1, p_2, p_3)\varphi_c(s) := (\varphi_c(s)^{-1}p_1, \varphi_c(s)^{-1}p_2, \varphi_c(s)^{-1}p_3)$. As it is easy to see, the lift of c that begins at another point $gp \in \pi^{-1}(c(a))$, $g \in \text{PU}V$, provides the same φ_c .

Let $c' : [b, d] \rightarrow S$ be one more piecewise vertical-horizontal path. Then, of course, $\varphi_{c \cup c'}(d) = \varphi_c(b) \circ \varphi_{c'}(d)$. In this way, we can explicitly find the paths φ_c 's for piecewise vertical-horizontal c 's if we know φ_c in the simple cases of a vertical/horizontal c .

Pick an arbitrary *basic* point $q_0 \in S$ and consider closed piecewise vertical-horizontal paths c that begin at q_0 . The corresponding φ_c 's form a subgroup $H_0(q_0) \leq \text{PGL}V$ called *rectangle holonomy group* at q_0 . For any $p \in \pi^{-1}(q_0)$, denote $H_0(p) := \{g \in \text{PU}V \mid gp = ph \text{ for some } h \in H_0(q_0)\} \leq \text{PU}V$. Since the lift of any vertical-horizontal path lies entirely in some E_F , we obtain $H_0(p) \leq C(F_p)$.

The rectangle holonomy group $H_0(q'_0)$ at another basic point $q'_0 \in S$ is isomorphic to $H_0(q_0)$ by means of the conjugation by φ_c , where c is a piecewise vertical-horizontal path from q_0 to q'_0 , $H_0(q_0) = \varphi_c H_0(q'_0) \varphi_c^{-1}$. As $H_0(q_0) \simeq H_0(p) \leq C(F_p)$ is commutative by Remark 3.9, this isomorphism is independent of the choice of c .

We can contract any closed piecewise vertical-horizontal path based at q_0 , keeping it piecewise vertical-horizontal and based at q_0 during the contraction. Therefore, $H_0(q_0)$ is path-connected. Any piecewise smooth closed path based at $q_0 \in S \setminus C$ can be approximated by a piecewise vertical-horizontal one. So, $H_0(q_0)$ is dense in $H(q_0)$. Since $H(q_0)$ is known to be a connected Lie group, we conclude that $H(p)$ is a closed connected subgroup in $C(F_p)$ for any $p \in \pi^{-1}(q_0)$. Moreover, $H(q_0) = H_0(q_0)$. Indeed, as $H(q_0)$ is abelian and $H_0(q_0)$ is path-connected, the subgroup $H_0(q_0)$ is the image of a linear subspace under the exponential map. It remains to observe that a small piecewise smooth closed path based at $q_0 \in S \setminus C$ can be approximated by a small piecewise vertical-horizontal one.

Let p be a real strongly regular triple and let R denote the corresponding real plane. Then $H_0(p) \leq \text{Stab}R$ because the bendings involving $p_1, p_2/p_2, p_3$ keep the triple inside R . Since $H_0(p)$ is path-connected, $H_0(p) \leq \text{Stab}^+ R$, where $\text{Stab}^+ R \leq \text{PU}V$ denotes the group of orientation-preserving isometries of the hyperbolic plane R . So, $H_0(p) \leq \text{Stab}^+ R \cap C(F_p)$. It is immediate that $\text{Stab}^+ R \cap C(F_p)$ is the 1-parameter subgroup generated by F_p in the group of isometries of the hyperbolic plane R .

Summarizing, we arrive at the following

4.2.1. Lemma. *The rectangle holonomy group coincides with the holonomy group, i.e., $H_0(p) = H(p)$ for any strongly regular triple $p \in E$. Moreover, $H_0(p)$ is a connected Lie subgroup in $C(F_p)$, $H_0(p) \leq C(F_p)$, where $F_p := R(p_3)R(p_2)R(p_1)$.*

If p is real, then $H_0(p)$ lies in the 1-parameter subgroup generated by F_p in the group of isometries of the hyperbolic plane R , $H_0(p) \leq \text{Stab}^+ R \cap C(F_p)$, where R stands for the real plane spanned by p ■

The main result of this paper is the following

4.2.2. Theorem. *Let $p := (p_1, p_2, p_3)$ be a strongly regular triple and let $F_p := R(p_3)R(p_2)R(p_1)$. Then $H_0(p) = C(F_p)$ unless the triple is real. If the triple is real, then $H_0(p) = \text{Stab}^+ R \cap C(F_p)$, where R stands for the real plane of the triple.*

4.2.3. Tangent space to E_F . In order to prove Theorem 4.2.2, we are going to explicitly find the curvature tensor of the connection on $E_F \rightarrow S$ and, using the Ambrose-Singer theorem, to calculate the holonomy group $H(p)$. So, we fix F .

A tangent vector $t \in T_p E_F$ can be described as a triple $t := (t_1, t_2, t_3)$ such that

$$(4.2.4) \quad \left. \frac{d}{d\varepsilon} \right|_{\varepsilon=0} R(p_3 + \varepsilon t_3(p_3)) R(p_2 + \varepsilon t_2(p_2)) R(p_1 + \varepsilon t_1(p_1)) = 0,$$

where $p = (p_1, p_2, p_3)$ and $t_i \in T_{p_i} \mathbb{P}_{\mathbb{C}} V \subset \text{Lin}_{\mathbb{C}}(V, V)$. One can easily verify that

$$\left. \frac{d}{d\varepsilon} \right|_{\varepsilon=0} R(p_i + \varepsilon t_i(p_i)) = 2(t_i + t_i^*) = 2\hat{t}_i R(p_i) \in T_{R(p_i)} \text{SU } V = \text{su } V \cdot R(p_i)$$

and that $\hat{t}_j R(p_j) + R(p_j) \hat{t}_j = 0$ (see Definition 2.3 for $\hat{t}$). Hence, (4.2.4) is equivalent to

$$\hat{t}_3 R(p_3) R(p_2) R(p_1) + R(p_3) \hat{t}_2 R(p_2) R(p_1) + R(p_3) R(p_2) \hat{t}_1 R(p_1) = 0,$$

i.e., to

$$(4.2.5) \quad -\hat{t}_3 + \hat{t}_2 + R(p_2) \hat{t}_1 R(p_2) = 0.$$

We have reduced the task of describing the tangent space $T_p E_F$ to the following one. Given a regular triple $p' = (p'_2, p'_1, p'_3)$ (where $p'_1 := R(p_2)p_1$, $p'_2 = p_2$, and $p'_3 = p_3$; note that $F = R(p'_3)R(p'_1)R(p'_2)$), we look for $t'_j \in T_{p'_j} \mathbb{P}_{\mathbb{C}} V$ (where $t'_1 := R(p_2)t_1$, $t'_2 := t_2$, and $t'_3 := -t_3$) such that $\hat{t}'_3 + \hat{t}'_2 + \hat{t}'_1 = 0$.

Let $[g'_{jk}]$ denote the Gram matrix of the p'_j 's and let $t'_j := \sum_k \langle -, p'_j \rangle a_{jk} p'_k$. The equation (4.2.5) is equivalent to the equalities $a_{kj} = \bar{a}_{jk}$ and the conditions $\sum_k a_{jk} p'_k \in p'^{\perp}_j$ are equivalent to the equalities $\sum_k a_{jk} g'_{kj} = 0$. They can be rewritten as $a_{jj} := \frac{1}{g'_{jj}} \text{Re} \left(\sum_{k \neq j} a_{jk} \right)$ and $\text{Im} \left(\sum_{k \neq j} a_{jk} g'_{kj} \right) = 0$ for all j . The last three equations in the a_{jk} 's, $j \neq k$, are linearly dependent (their sum equals zero) and define a 4-dimensional space of solutions because $g'_{jk} \neq 0$.

4.2.6. Lemma. *The vector fields*

$$b_1(p) := (\langle -, p_1 \rangle (\frac{p_1}{g_{11}} - \frac{p_2}{g_{21}}), \langle -, p_2 \rangle (\frac{p_1}{g_{12}} - \frac{p_2}{g_{22}}), 0), \quad b_2(p) := (0, \langle -, p_2 \rangle (\frac{p_2}{g_{22}} - \frac{p_3}{g_{32}}), \langle -, p_3 \rangle (\frac{p_2}{g_{23}} - \frac{p_3}{g_{33}}))$$

defined for all $p := (p_1, p_2, p_3) \in E$ never vanish and span the horizontal $\mathbb{R}$ -linear space of the connection on $E \rightarrow S$ over $S \setminus C$, where $[g_{jk}]$ stands for the Gram matrix of the p_j 's. Denote $\tau := \frac{g_{13}g_{22}}{g_{12}g_{23}}$. Then

$$[b_1, b_2](p) = \left(\langle -, p_1 \rangle \bar{\tau} (\frac{p_2}{g_{21}} - \frac{p_3}{g_{31}}), \langle -, p_2 \rangle ((2 - \tau) \frac{p_1}{g_{12}} + 2i \text{Im } \tau \frac{p_2}{g_{22}} + (\bar{\tau} - 2) \frac{p_3}{g_{32}}), \langle -, p_3 \rangle \tau (\frac{p_1}{g_{13}} - \frac{p_2}{g_{23}}) \right).$$

Proof. It is immediate that the vectors b_1 and b_2 never vanish and are tangent to the lifts of the vertical and horizontal lines, respectively.

Let f_j be an analytic function on E depending only on $p_j \in \mathbb{P}_{\mathbb{C}} V$, $j = 1, 2, 3$. Then

$$b_1 f_1 = \left. \frac{d}{d\varepsilon} \right|_{\varepsilon=0} f_1(p_1 - \varepsilon \frac{g_{11}}{g_{21}} p_2), \quad b_1 f_2 = \left. \frac{d}{d\varepsilon} \right|_{\varepsilon=0} f_2(p_2 + \varepsilon \frac{g_{22}}{g_{12}} p_1), \quad b_1 f_3 = 0,$$

$$b_2 f_1 = 0, \quad b_2 f_2 = \left. \frac{d}{d\varepsilon} \right|_{\varepsilon=0} f_2(p_2 - \varepsilon \frac{g_{22}}{g_{32}} p_3), \quad b_2 f_3 = \left. \frac{d}{d\varepsilon} \right|_{\varepsilon=0} f_3(p_3 + \varepsilon \frac{g_{33}}{g_{23}} p_2),$$

$$[b_1, b_2] f_1 = -b_2 b_1 f_1 = -\left. \frac{d}{d\delta} \right|_{\delta=0} \left. \frac{d}{d\varepsilon} \right|_{\varepsilon=0} f_1 \left(p_1 - \varepsilon g_{11} \frac{\frac{p_2}{g_{21}} + \delta \frac{g_{22}}{g_{21}} (\frac{p_2}{g_{22}} - \frac{p_3}{g_{32}})}{1 + \delta \frac{g_{22}}{g_{21}} (\frac{g_{21}}{g_{22}} - \frac{g_{31}}{g_{32}})} \right) = \langle -, p_1 \rangle \frac{g_{22}g_{31}}{g_{32}g_{21}} (\frac{p_2}{g_{21}} - \frac{p_3}{g_{31}}) f_1.$$

Similarly, $[b_1, b_2] f_3 = \langle -, p_3 \rangle \tau (\frac{p_1}{g_{13}} - \frac{p_2}{g_{23}}) f_3$.

We have calculated the first and the last components of the vector $[b_1, b_2](p)$. Since the vectors $b_1(p')$, $b_2(p')$ are tangent to E_{F_p} for all $p' \in E_{F_p}$, the vector $[b_1, b_2](p)$ has to be tangent to E_{F_p} . It is straightforward that the right-hand side of the expression given in Lemma 4.2.6 satisfies the equation (4.2.5), which uniquely determines the second component ■

4.2.7. Curvature tensor of $E_F \rightarrow S$. Let $l \in \text{su } V$. By [AGG, Lemma 4.1.4], the associated to l tangent vector to the fibre of $\pi : E \rightarrow S$ at $p := (p_1, p_2, p_3) \in E$ has the form

$$(\langle -, p_1 \rangle \frac{\pi[p_1]l(p_1)}{g_{11}}, \langle -, p_2 \rangle \frac{\pi[p_2]l(p_2)}{g_{22}}, \langle -, p_3 \rangle \frac{\pi[p_3]l(p_3)}{g_{33}}),$$

where $\pi[p_j] : V \rightarrow p_j^\perp$ denotes the orthogonal projection and $[g_{jk}]$ is the Gram matrix of the p_j 's. Conversely, let $t := (\langle -, p_1 \rangle q_1, \langle -, p_2 \rangle q_2, \langle -, p_3 \rangle q_3)$ be a tangent vector to E at $p := (p_1, p_2, p_3) \in E$ with $q_j \in p_j^\perp$. Denote by T the $\mathbb{C}$ -linear map $T : V \rightarrow V$ given by $T(p_j) := g_{jj}q_j$ and consider a $\mathbb{C}$ -linear map $L : V \rightarrow V$ such that $L(p_j) := T(p_j) - id_j p_j$ with $d_j \in \mathbb{R}$ for all $j = 1, 2, 3$. Then t is tangent to the fibre iff $L \in \text{su } V$ for suitable d_j 's. In terms of the basis p_1, p_2, p_3 , this means that $\text{tr } T = i \text{tr } D$ and $T^t[g_{jk}] + [g_{jk}]\overline{T} = i(D[g_{jk}] - [g_{jk}]D)$, where $D := \begin{bmatrix} d_1 & 0 & 0 \\ 0 & d_2 & 0 \\ 0 & 0 & d_3 \end{bmatrix}$. In other words, t is tangent to the fibre of $\pi : E \rightarrow S$ iff $\text{tr } T \in i\mathbb{R}$ and

$$(4.2.8) \quad T^t[g_{jk}] + [g_{jk}]\overline{T} = i \begin{bmatrix} 0 & a_{12}g_{12} & a_{13}g_{13} \\ a_{21}g_{21} & 0 & a_{23}g_{23} \\ a_{31}g_{31} & a_{32}g_{32} & 0 \end{bmatrix}$$

with $a_{kj} = -a_{jk} \in \mathbb{R}$ and $a_{12} + a_{23} + a_{31} = 0$. Moreover, in this case, $d_1 = \frac{d+a_{12}-a_{31}}{3}$, $d_2 = \frac{d+a_{23}-a_{12}}{3}$, $d_3 = \frac{d+a_{31}-a_{23}}{3}$, where $id := \text{tr } T$. In particular,

$$(4.2.9) \quad d_1 = \frac{d+2a_{12}+a_{23}}{3}.$$

Denote by ω and Ω the 1-form of the connection on $E_F \rightarrow S$ and its curvature tensor, both defined over $S \setminus C$. They take values in the Lie algebra $\mathfrak{c}(F) \leq \text{su } V$ of the group $C(F)$. Since $\Omega(b_2 \wedge b_1) = \omega[b_1, b_2]$ by Lemma 4.2.6, we need the following lemma.

4.2.10. Lemma. *Let $p := (p_1, p_2, p_3) \in E_F \setminus \pi^{-1}(C)$. Then*

$$\omega[b_1, b_2](p_1) = g_{11} \frac{(1-\beta-t_2)t_1 t_2 - 2\alpha^2}{t_1^2 t_2^2 (t-1)} \left(\frac{p_1}{g_{11}} - \frac{p_2}{g_{21}} \right) + g_{11} \overline{\tau} \left(\frac{p_2}{g_{21}} - \frac{p_3}{g_{31}} \right) + \frac{i\alpha(1-\beta-3t_2+2t_1 t_2)}{3t_1^2 t_2^2 (t-1)} p_1,$$

where $\tau := \frac{g_{13}g_{22}}{g_{12}g_{23}}$ and $[g_{jk}]$ stands for the Gram matrix of the p_j 's.

Proof. By Lemma 4.2.6, there exist unique $c_1, c_2 \in \mathbb{R}$ such that the vector $c_1 b_1(p) + c_2 b_2(p) + [b_1, b_2](p)$ is tangent to the fibre $E_F \cap \pi^{-1}(q)$ of $E_F \rightarrow S$. By Lemma 4.2.6 and the considerations in 4.2.7, $\omega[b_1, b_2](p_1) = g_{11}c_1 \left(\frac{p_1}{g_{11}} - \frac{p_2}{g_{21}} \right) + g_{11}\overline{\tau} \left(\frac{p_2}{g_{21}} - \frac{p_3}{g_{31}} \right) - id_1 p_1$.

The matrices of $b_1(p), b_2(p), [b_1, b_2](p)$ are

$$B_1 := \begin{bmatrix} 1 & \frac{g_{22}}{g_{12}} & 0 \\ -\frac{g_{11}}{g_{21}} & -1 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad B_2 := \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & \frac{g_{33}}{g_{23}} \\ 0 & -\frac{g_{22}}{g_{32}} & -1 \end{bmatrix}, \quad B := \begin{bmatrix} 0 & \frac{g_{22}}{g_{12}}(2-\tau) & \frac{g_{33}}{g_{13}}\tau \\ \frac{g_{11}}{g_{21}}\overline{\tau} & 2i \text{Im } \tau & -\frac{g_{33}}{g_{23}}\tau \\ -\frac{g_{11}}{g_{31}}\overline{\tau} & \frac{g_{22}}{g_{32}}(\overline{\tau}-2) & 0 \end{bmatrix}.$$

We calculate the hermitian matrices

$$B_1^t[g_{jk}] + [g_{jk}]\overline{B}_1 = \begin{bmatrix} 0 & 0 & * \\ 0 & 0 & g_{23}(\tau-1) \\ g_{31}(1-\frac{g_{11}g_{32}}{g_{31}g_{12}}) & * & 0 \end{bmatrix}, \quad B_2^t[g_{jk}] + [g_{jk}]\overline{B}_2 = \begin{bmatrix} 0 & g_{12}(1-\tau) & * \\ * & 0 & 0 \\ g_{31}(\frac{g_{21}g_{33}}{g_{23}g_{31}}-1) & 0 & 0 \end{bmatrix},$$

$$B^t[g_{jk}] + [g_{jk}]\overline{B} = \begin{bmatrix} 0 & g_{12}(\frac{g_{11}g_{22}}{g_{12}g_{21}} + \tau(\tau-3) + \overline{\tau}) & * \\ * & 0 & -g_{23}(\frac{g_{22}g_{33}}{g_{23}g_{32}} + \tau(\tau-3) + \overline{\tau}) \\ g_{31}(\frac{g_{13}g_{22}}{g_{12}g_{23}}(\frac{g_{11}g_{32}}{g_{31}g_{12}} - \frac{g_{21}g_{33}}{g_{23}g_{31}})) & * & 0 \end{bmatrix}.$$

Let $T := c_1 B_1 + c_2 B_2 + B$. Then $\text{tr } T = id$, where $d := 2t'$ and $t + it' := \tau$.

The 12- and 23-components of the matrix $T^t[g_{jk}] + [g_{jk}]\overline{T}$ belong to $ig_{12}\mathbb{R}$ and $ig_{12}\mathbb{R}$ iff

$$c_2(t-1) = \text{Re}\left(\frac{g_{11}g_{22}}{g_{12}g_{21}} + \tau(\tau-3) + \overline{\tau}\right), \quad c_1(t-1) = \text{Re}\left(\frac{g_{22}g_{33}}{g_{23}g_{32}} + \tau(\tau-3) + \overline{\tau}\right),$$

respectively. Moreover, we can also see that

$$a_{12} = \text{Im}(-c_2\tau + \tau(\tau-3) + \overline{\tau}), \quad a_{23} = \text{Im}(c_1\tau - \tau(\tau-3) - \overline{\tau}).$$

Hence,

$$c_1 = \frac{\frac{1}{t_2} + t^2 - t'^2 - 2t}{t-1}, \quad c_2 = \frac{\frac{1}{t_1} + t^2 - t'^2 - 2t}{t-1}, \quad a_{12} = -c_2 t' + 2t t' - 4t', \quad a_{23} = c_1 t' - 2t t' + 4t'.$$

By (4.2.9), $d_1 = \frac{(c_1 - 2c_2 + 2t - 2)t'}{3} = \frac{(\frac{1}{t_2} - \frac{2}{t_1} + t^2 + t'^2 - 2t + 2)t'}{3(t-1)}$.

From $t_1 t_2 \overline{\tau} = \frac{g_{12}g_{23}g_{31}}{g_{11}g_{22}g_{33}}$, we conclude that $t' = -\frac{\alpha}{t_1 t_2}$. Using equation (4.1.2), we can see that $c_1 = \frac{(1-\beta-t_2)t_1 t_2 - 2\alpha^2}{t_1^2 t_2^2 (t-1)}$ and $d_1 = -\frac{\alpha(1-\beta-3t_2+2t_1 t_2)}{3t_1^2 t_2^2 (t-1)}$. ■

Proof of Theorem 4.2.2. By Lemma 4.2.10, the holonomy group is not trivial. So, we assume that $\alpha \neq 0$.

We fix the point p_1 and the geodesic $G \setminus \langle p_2, p_3 \rangle$. It suffices to show that, for a generic $(p_1, p_2, p_3) \in E_F$, the element $\omega[b_1, b_2](p_1) \in V$ does not remain $\mathbb{R}$ -proportional when we bend p_2, p_3 .

Let $\{u_3\} := \mathbb{P}_{\mathbb{C}} p_1^\perp \cap L(p_2, p_3)$ and let u_2 be the polar point to $L(p_1, u_3)$. Then $\frac{\langle p_2, u_2 \rangle}{g_{21}} = \frac{\langle p_3, u_2 \rangle}{g_{31}}$ because $\frac{p_2}{g_{21}} - \frac{p_3}{g_{31}} \in \mathbb{P}_{\mathbb{C}} p_1^\perp \cap L(p_2, p_3)$. Note that $b := \frac{\langle g, u_2 \rangle}{\langle g, p_1 \rangle}$ is independent of the choice of $g \in G \setminus \langle p_2, p_3 \rangle \setminus \mathbb{P}_{\mathbb{C}} p_1^\perp$. Indeed, $\frac{\langle g, u_2 \rangle}{\langle g, p_1 \rangle} = \frac{(1 + \frac{r g_{31}}{g_{32} g_{21}}) \langle p_2, u_2 \rangle}{g_{21} + \frac{r g_{31}}{g_{32}}} = \frac{\langle p_2, u_2 \rangle}{g_{21}}$ because $\langle p_3, u_2 \rangle = \frac{g_{31} \langle p_2, u_2 \rangle}{g_{21}}$, where $g := p_2 + \frac{r p_3}{g_{32}} \in G \setminus \langle p_2, p_3 \rangle$ for any $r \in \mathbb{R}$. The point u_2 cannot be polar to $L(p_2, p_3)$. Therefore, $b \neq 0$.

It remains to observe that

$$\langle \omega[b_1, b_2](p_1), u_2 \rangle = \frac{2\alpha^2 - (1-\beta-t_2)t_1 t_2}{t_1^2 t_2^2 (t-1)} g_{11} b, \quad \langle \omega[b_1, b_2](p_1), p_1 \rangle = \frac{i\alpha(1-\beta-3t_2+2t_1 t_2)}{3t_1^2 t_2^2 (t-1)} g_{11}$$

by Lemma 4.2.10. ■

4.3. Regular triples. In this subsection, we discuss regular triples that are not strongly regular and, a little bit, spherical configurations.

Consider a regular triple $p_1, p_2, p_3 \in L$ that lies in a projective line L . The case when all points are negative is in fact covered by Theorem 4.2.2 : it does not differ from the case of a real strongly regular triple because the geometries of the Poincaré disc and of the Beltrami-Klein disc are essentially the same. When a point p_j is positive, we can substitute p_j by the negative point $p'_j \in L$ orthogonal to p_j . Indeed, let p denote the point polar to L . Then $R(p)$ commutes with every $R(p_k)$ and $R(p)R(p_j) = R(p'_j)$. Every geodesic that passes through p_j passes necessarily through p'_j [AGr, Section 3]. Now, it is easy to see that a bending involving p_j, p_k corresponds to a bending involving p'_j, p_k .

Let $p_1, p_2, p_3 \in R$ be a real regular triple with a positive p_j , where R stands for the real plane spanned by the p_k 's. In terms of the Beltrami-Klein disc D of all negative points in R , the set $G := D \cap \mathbb{P}_{\mathbb{C}} p_j^\perp$ is a geodesic. Denote by Γ the geodesic in D that passes through p_k and is perpendicular to G . When we bend p_j, p_k , we in fact move G and p_k keeping the distance between G and p_k , keeping G orthogonal to Γ , and keeping $p_k \in \Gamma$. At some moment, the third point p_l lies in G , $p_l \in G$. If necessary, we can replace $R(p_j)$ and $R(p_k)$ by $R(R(p_j)p_k)$ and $R(p_j)$. Now, two subsequent points p_j, p_l are orthogonal and $R(p_j)R(p_l)$ equals some $R(p)$ due to orthogonal relations.

If one discards the requirement that at most one point is positive, there is an example of a triple p_1, p_2, p_3 such that the projective lines $L(p_1, p_2), L(p_2, p_3)$ are hyperbolic and $L(p_1, p_3)$ becomes spherical after a bending involving p_1, p_2 . Indeed, there is a number $z \in \mathbb{C}$ such that $|\frac{1}{4} + z| < \frac{1}{2}$ and $1 < |1 + z|$ (for instance, $z = \frac{1}{8}$). By Sylvester's Criterion, there are points $v_1, v_2, p_3 \in V$ with the Gram matrix $\begin{bmatrix} 0 & \frac{1}{2} & 1 \\ \frac{1}{2} & 0 & \bar{z} \\ 1 & z & 1 \end{bmatrix}$. We put

$$p_1 := 2v_1 - \frac{1}{2}v_2, \quad p_2 := v_1 + v_2, \quad p'_1 := v_1 - v_2, \quad p'_2 := \frac{1}{2}v_1 + 2v_2.$$

Then the projective lines $L(p_1, p_2), L(p_2, p_3)$ are hyperbolic because $1 < |1 + z|$, whereas the projective line $L(p'_2, p_3)$ is spherical because $|\frac{1}{2} + 2z| < 1$.

Nevertheless, in principle, there can exist a *spherical* configuration of points $p_1, p_2, \dots, p_n$, i.e., a non-trivial configuration subject to the relation $R(p_n) \dots R(p_2)R(p_1) = 1$ with spherical projective lines $L(p_{j-1}, p_j)$ (the indices are modulo n) that remain spherical after a finite number of bendings. In this case, one should suitably modify Conjecture 1.1.

While looking for basic relations, one should not go too far. Most likely, every isometry in $SU V$ is a product of 5 reflections. Hence, it must be sufficient to deal with $n \leq 11$. However, it is quite possible that actually we only need to study the relations with $n \leq 6$.

5. Pentagons

In this section, we begin to study 3 basic relations (in the sense of Conjecture 1.1), the pentagons. (Yet, we do not know if there exists a spherical configuration with $n = 5$.) A *pentagon* is a configuration $p_1, p_2, p_3, p_4, p_5 \in \mathbb{P}_{\mathbb{C}} V \setminus S V$ of 5 nonisotropic points such that at most one of the p_j 's is positive, p_{j-1} is not orthogonal nor equal to p_j for all j (the indices are modulo 5), and

$$R(p_5)R(p_4)R(p_3)R(p_2)R(p_1) = \delta$$

in $SU V$, where $\delta^3 = 1$. We will show that pentagons considered modulo congruence form exactly 3 components. Moreover, two pentagons with the same δ are connected by means of a finite number of bendings. Each component is a smooth 4-manifold.

If $\delta = 1$, the pentagons are known to provide faithful and discrete representations $\varrho : H_5 \rightarrow PU V$ [ABG], where H_n denotes the *hyperelliptic* group, i.e., the group generated by $r_1, r_2, \dots, r_n$ with the defining relations $r_j^2 = 1$, $j = 1, 2, \dots, n$, and $r_n \dots r_2 r_1 = 1$. Moreover, in this case the p_j 's are all negative and span a real plane R . In other words, the representation ϱ is $\mathbb{R}$ -fuchsian as R is ϱH_5 -stable. So, we arrive at a sort of Toledo rigidity [Tol] : while deforming a pentagon with $\delta = 1$, the representation ϱ remains faithful, discrete, and $\mathbb{R}$ -fuchsian.

The two components with $\delta \neq 1$ are congruent: one can be obtained from the other by reflecting the pentagons in a real plane (they are 'complex conjugated'). One of the p_j 's is always positive. In [Ana], we construct a pentagon with $\delta \neq 1$ that provides a discrete and faithful representation $\varrho : H_5 \rightarrow PU V$. We believe that every pentagon provides a faithful and discrete representation (Conjecture 1.2).

5.1. Signs of points. It follows from $R(p_3)R(p_2)R(p_1) = \delta R(p_4)R(p_5)$ and Lemma 3.1 that

$$(5.2) \quad 8i\alpha(p_1, p_2, p_3) + 4\beta(p_1, p_2, p_3) - 1 = \delta(4\text{ta}(p_4, p_5) - 1).$$

Suppose that $\delta = 1$ and $p_1, p_2, p_3 \in BV$. Then $\alpha(p_1, p_2, p_3) = 0$, $\beta(p_1, p_2, p_3) \geq 0$, and $\beta(p_1, p_2, p_3) = \text{ta}(p_4, p_5)$, implying that p_4, p_5 have the same sign, i.e., are negative.

Suppose that $\delta \neq 1$ and $p_4, p_5 \in BV$. Then $\text{Re } \delta = -\frac{1}{2}$ and $4\beta(p_1, p_2, p_3) - 1 = -\frac{1}{2}(4\text{ta}(p_4, p_5) - 1)$. Hence, $8\beta(p_1, p_2, p_3) = 3 - 4\text{ta}(p_4, p_5)$. Since $\text{ta}(p_4, p_5) > 1$, we obtain $\beta(p_1, p_2, p_3) < 0$. So, one of the points p_1, p_2, p_3 is positive.

5.3. Existence. For distinct $p_4, p_5 \in BV$, the eigenvectors of $R(p_4)R(p_5)$ are the vertices of the geodesic $G\langle p_4, p_5 \rangle$ and the point polar to $L(p_4, p_5)$. The corresponding eigenvalues are $s^{-1}, s, 1$, where $s > 1$. By Proposition 3.8, we can find a regular triple p_1, p_2, p_3 such that $R(p_3)R(p_2)R(p_1) = \delta R(p_4)R(p_5)$. If $p_3 = p_4$ or $p_1 = p_5$, then p_1, p_2 or p_2, p_3 are orthogonal by Remark 3.2, which contradicts the fact that the triple p_1, p_2, p_3 is regular. If p_1, p_5 are orthogonal, then, after applying the orthogonal relation, we obtain $R(p_4)R(p_3)R(p_2)R(p_1) = \delta$ with p, p_1, p_5 pairwise orthogonal. By Corollary 3.3, p, p_2, p_3, p_4 lie on the geodesic $G\langle p_2, p_3 \rangle$. As p, p_1 are orthogonal, $p_1 \in G\langle p_2, p_3 \rangle$, which contradicts the fact that the triple p_1, p_2, p_3 is regular. Symmetric arguments work if p_3, p_4 are orthogonal.

Thus, we have shown that, for any δ with $\delta^3 = 1$, there exists a pentagon.

5.4. Connectedness by means of bendings. Given a pentagon, the triple p_1, p_2, p_3 is strongly regular. Indeed, if p_1, p_2, p_3 are on a same geodesic G , then $R(p_3)R(p_2)R(p_1) = R(p)$ with $p \in G$ by Corollary 3.3. Now, $R(p_5)R(p_4)R(p) = \delta$ implies that p_4, p_5 are orthogonal by Remark 3.2. A contradiction. If p_1, p_2, p_3 are on a same projective line, then $\beta(p_1, p_2, p_3) = 0$ and (5.2) implies that $\operatorname{Re} \delta = -\frac{1}{2}$ and, hence, $4 \operatorname{ta}(p_4, p_5) = 3$. So, the projective line $L(p_4, p_5)$ is spherical; a contradiction. If p_1, p_2, p_3 are on a same real plane, then $\alpha(p_1, p_2, p_3) = 0$. Therefore, $\delta = 1$ by (5.2) and the p_j 's are all negative by 5.1.

In particular, after any bending, a pentagon remains a pentagon. Indeed, after a bending involving, say, p_2, p_3 , the triples p_1, p_2, p_3 and p_2, p_3, p_4 remain strongly regular by Corollary 4.1.5.

Given two pentagons with the same δ and negative p_2, p_3, p_4, p_5 , by means of a bending involving p_3, p_4 , one can make $\operatorname{ta}(p_4, p_5)$ arbitrarily big. So, we can assume that $\operatorname{ta}(p_4, p_5)$ is the same for both pentagons. The connectedness follows now from Corollary 4.1.5.

5.5. Real pentagons. Consider the case of $\delta = 1$. By 5.1 and 5.4, the points p_1, p_2, p_3 form a strongly regular real triple and, therefore, span a real plane R , which is stable under the isometry $I := R(p_3)R(p_2)R(p_1) = R(p_4)R(p_5)$. As the fixed points of the isometry $R(p_4)R(p_5)$ are the (isotropic) vertices v, v' of the geodesic $G\langle p_4, p_5 \rangle$ and the (positive) point polar to the projective line $L(p_4, p_5)$, the isometry I of the Beltrami-Klein disc $D \subset R$ has to be hyperbolic. So, its isotropic fixed points should coincide with v, v' , implying $v, v' \in R$ and, hence, $p_4, p_5 \in R$. As there is no essential difference between the Beltrami-Klein disc and the Poincaré one, we can apply [ABG, Corollary 3.16] and conclude that the pentagon provides a faithful and discrete representation $\varrho : H_5 \rightarrow \operatorname{PU} V$.

5.6. Smoothness. The space of real pentagons is smooth and 4-dimensional by [ABG, Corollary 3.17]. So, we assume $\delta := -\frac{1}{2} + i\frac{\sqrt{3}}{2}$ and p_1 positive.

Let $t_j := \operatorname{ta}(p_j, p_{j+1})$ (the indices are modulo 5). It follows from (5.2) that $\alpha(p_1, p_2, p_3) = \frac{\sqrt{3}(4t_4-1)}{16}$ and $\beta(p_1, p_2, p_3) = \frac{3-4t_4}{8}$. The inequalities (4.1.3) for t_1, t_2, t_4 are

$$(5.7) \quad t_1 < 0, \quad 1 < t_2, \quad 1 < t_4.$$

Therefore, $\beta < 0$, as desired. The equation (4.1.2) takes the form

$$(5.8) \quad (t_1 - 1)(t_2 - 1) = t_1 t_2 (t - 1)^2 + \frac{3(4t_4-1)^2}{256t_1 t_2} + \frac{3-4t_4}{8}.$$

The equation (5.8) in t_1, t_2, t_4, t together with the inequalities (5.7) defines a smooth 3-manifold $T \subset \mathbb{R}^4(t_1, t_2, t_4, t)$ fibred by Lemma 4.1.1 into planes over the axis $A := \{t_4 \mid t_4 > 1\}$. Hence, $T \simeq \mathbb{R}^3$. Geometrically, a pentagon is determined by a point in T uniquely up to a bending involving p_4, p_5 . In other words, the space of pentagons with a given δ is diffeomorphic to $\mathbb{R}^4$.

6. References

- [ABG] S. Anan'in, E. C. B. Gonçalves, *A hyperelliptic view on Teichmüller space*. I, preprint <http://arxiv.org/abs/0709.1711>
- [AGG] S. Anan'in, C. H. Grossi, N. Gusevskii, *Complex hyperbolic structures on disc bundles over surfaces*, Int. Math. Res. Not. **2011** (2011), no. 19, 4295–4375, see also <http://arxiv.org/abs/math/0511741>
- [AGr] S. Anan'in, C. H. Grossi, *Coordinate-free classic geometries*, Mosc. Math. J. **11** (2011), 633–655, see also <http://arxiv.org/abs/math/0702714>
- [Ana] S. Anan'in, *A discrete pentagon*, in preparation
- [Gol1] W. M. Goldman, *Topological components of spaces of representations*, Invent. Math. **93** (1988), no. 3, 557–607
- [Gol2] W. M. Goldman, *Complex hyperbolic geometry*, Oxford Mathematical Monographs. Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1999, xx+316 pp.
- [Pra] A. Pratoussevitch, *Traces in complex hyperbolic triangle groups*, Geometriae Dedicata **111** (2005), 159–185
- [Tol] D. Toledo, *Representations of surface groups in complex hyperbolic space*, J. Differential Geom. **29** (1989), no. 1, 125–133

DEPARTAMENTO DE MATEMÁTICA, IMECC, UNIVERSIDADE ESTADUAL DE CAMPINAS,
13083-970—CAMPINAS—SP, BRASIL